Download the WhaleReport app | wildwhales.org | 1.866.J.SAW.ONE





Harbour porpoise

Phocoena phocoena

LENGTH Less than 2 m / 6 ft
DORSAL FIN Dark, triangular
APPEARANCE Dark grey back with lighter underside
BEHAVIOUR Inconspicuous, travels slowly;
generally alone or in groups of 2–3





Dall's porpoise

Phocoenoides dalli

LENGTH Up to 2.2 m / 7 ft
DORSAL FIN Triangular with white trailing edge
APPEARANCE Black, with large white flanks; small hump on back
before tail flukes
BEHAVIOUR Usually in groups of 2–10; creates "rooster-tail" of
spray when swimming quickly






Pacific white-sided dolphin

Lagenorhynchus obliquidens

LENGTH Up to 2.5 m / 8 ft
DORSAL FIN Sharply curved, bi-coloured
APPEARANCE Dark back, white underside, grey side streaks
BEHAVIOUR Only small, local cetacean that leaps completely
clear of water; usually in groups of 50+





This project was undertaken with the financial support of the Government of Canada.

Ce projet a été réalisé avec l'appui financier

Ce projet a été réalisé avec l'appui financier du gouvernement du Canada.



Killer whale

Orcinus orca

LENGTH 6-9 m / 18-28 ft
DORSAL FIN Black, triangular in females and calves;
tall, distinctive in adult males
APPEARANCE Black body with white underside;
grey saddle patch behind dorsal fin; white eye patch
BEHAVIOUR Usually in groups of 3-25;
bushy blows visible in cold weather

surfaces 1-2 times between dives: elusive

Download the WhaleReport app | wildwhales.org | 1.866.I.SAW.ONE





Minke whale

Balaenoptera acutorostrata

LENGTH 7-10 m / 23-33 ft

DORSAL FIN Curved, far back on body

APPEARANCE Dark grey to black colouration; white band on topside
of pectoral fins; pointed head with prominent noseridge

BEHAVIOUR Usually solitary; blows rarely visible;





Grey whale

Eschrichtius robustus

LENGTH 12-14 m / 39-46 ft.

DORSAL FIN None; knuckle-like bumps on lower back

APPEARANCE Mottled grey colour; patches of barnacles

BEHAVIOUR Low, bushy, heart- or v-shaped blow; occasionally

lifts tail flukes when diving; often feeds close to shore





Humpback whale

Megaptera novaeangliae

LENGTH 11–15 m / 37–50 ft

DORSAL FIN Small, knobby fin with broad base

APPEARANCE Black body; bumps or knobs on head;
long pectoral fins with white underside

BEHAVIOUR Medium-sized, bushy blow; usually surfaces 2-5 times between deep dives; often raises tail flukes on final dive





This project was undertaken with the financial support of the Government of Canada.

Ce projet a été réalisé avec l'appui financier du gouvernement du Canada.



Fin whale

Balaenoptera physalus

LENGTH 18-22 m / 59-72 ft

DORSAL FIN Sickle-shaped; relatively large; very far back on body APPEARANCE Right lower lip is white, while left is dark;

Blowhole and dorsal fin not visible simultaneously when surfacing BEHAVIOUR Tall, narrow blow; rarely lifts tail flukes when diving



Download the WhaleReport app | wildwhales.org | 1.866.I.SAW.ONE





Sperm whale

Physeter macrocephalus

LENGTH 11-18 m / 36-59 ft DORSAL FIN Triangular/rounded hump, followed by knuckles APPEARANCE Wrinkled dark skin; huge square-shaped head; blowhole very close to front of head BEHAVIOUR Low, bushy blow, angled to the left; lifts broad triangular tail flukes high in air before diving





Sei whale

Balaenoptera borealis

LENGTH 12-14 m / 39-46 ft DORSAL FIN Slender and erect; less than 2/3 from front of body APPEARANCE Blue-grey, dark grey or black back; blowhole and dorsal fin visible simultaneously when surfacing BEHAVIOUR Narrow, cloud-like blow; rarely exposes tail flukes





Blue whale

Balaenoptera musculus

LENGTH 20-25 m / 65-83 ft DORSAL FIN Variable shape; very small relative to body APPEARANCE Blue-grey, mottled colouring; dorsal fin not seen until well after blow ends BEHAVIOUR Blows 9 m / 30 ft tall; often exposes tail flukes when diving





This project was undertaken with the financial support of the Government of Canada.

Ce projet a été réalisé avec l'appui financier

North Pacific right whale

Eubalaena japonica

LENGTH 11-15 m / 37-50 ft DORSAL FIN None

APPEARANCE Dark, rotund body; large head covered in white, irregular growths (callosities)

BEHAVIOUR Wide, V-shaped blow; may raise tail flukes when diving

NOTE: North Pacific right whales have not been seen in B.C. waters in 40 years. If you see one, please call 1.866.I.SAW.ONE





Sea turtles of B.C. identification guide

Download the WhaleReport app | wildwhales.org | 1.866.I.SAW.ONE





Leatherback sea turtle

Dermochelys coriacea (The most commonly seen sea turtle in B.C. waters)

TOTAL LENGTH Up to 3 m / 9 ft COLOUR Black, with white spots FEATURES Very large; ridges along back (looks like the bottom of a boat); shell tapers to a point at rear end








Green sea turtle

Chelonia mydas

TOTAL LENGTH Up to 1.5 m / 5 ft COLOUR Dark greyish green with light blotches on shell FEATURES Shell tapered at rear; small head relative to body; four costal (side) plates (see • on diagram)








Loggerhead turtle

Caretta caretta

TOTAL LENGTH 0.3-1 m / 1-3 ft (juvenile only) COLOUR Reddish brown FEATURES Large head and jaws; five costal (side) plates (see • on diagram)







of the Government of Canada.


Ce projet a été réalisé avec l'appui financier



Olive ridley turtle Lepidochelys olivacea

TOTAL LENGTH Up to 1 m / 3 ft **COLOUR** Olive/grey FEATURES Shell width equal to or greater than shell length (highly domed shell); six to nine costal (side) plates (see • on diagram)

NOTE: Loggerhead and olive ridley turtles have not yet been recorded in B.C. waters but have been seen in Washington and Alaska.







Download the WhaleReport app | wildwhales.org | 1.866.I.SAW.ONE





Northern right whale dolphin

Lissodelphis borealis

LENGTH Up to 3 m / 9 ft
DORSAL FIN None, only dolphin species in the North Pacific without
APPEARANCE Slender bodies; mostly black with white on the belly
and the tip of the rostrum (beak); very small pectoral flippers
BEHAVIOUR Extremely social, usually in large groups of 200+;
often leap clear of the water; may associate with other dolphin species





Risso's dolphin

Grampus griseus

LENGTH Up to 4 m / 12 ft
DORSAL FIN Tall, curved; usually darker grey than body
APPEARANCE Grey; body often appears covered in scratches;
head is very rounded without a noticeable rostrum (beak)
BEHAVIOUR Social, usually in groups of 10-50; may porpoise, breach, spyhop





Common dolphin

Delpinus delphis

LENGTH Up to 2.5 m / 8 ft

DORSAL FIN Small, highly pointed and black

APPEARANCE Criss-cross pattern of yellow, white, grey and black bands

(note: animal in photo is covered in diatoms, giving a more brownish colouration);

prominent, pointy black rostrum (beak)

BEHAVIOUR Very acrobatic: social often in groups of 20-100+:

BEHAVIOUR Very acrobatic; social, often in groups of 20-100+; in B.C. small groups (2-3) or lone animals have been observed





This project was undertaken with the financial support of the Government of Canada.

Ce projet a été réalisé avec l'appui financier



False killer whale

Pseudorca crassidens

LENGTH Up to 6 m / 18 ft
DORSAL FIN Curved, black
APPEARANCE All black body; small, rounded head; long, slightly bent pectoral flippers
BEHAVIOUR Extremely social, usually in large groups of 20-100;

in B.C. lone animals have been observed; may breach, bowride, or porpoise



Download the WhaleReport app | wildwhales.org | 1.866.J.SAW.ONE



Beaked whales are found offshore in deep waters. There are many species of beaked whale in the northern Pacific Ocean and this sheet includes the most frequently encountered. Beaked whales avoid vessels and are usually found far from shore; few sighting reports of beaked whales exist in British Columbia.

BEAKED WHALE IDENTIFICATION: Single blowhole. Most individuals (especially males) are heavily scarred. Beaked whales have prominent rostrums (beaks) and teeth that emerge from their lower jaw (primarily males).

BEAKED WHALE BEHAVIOUR: Beaked whales spend very little time at the surface and are deep divers. Groups may range from ~5-20 and individuals are often seen swimming in very close proximity to one another.

Baird's beaked whale

Berardius bairdii

LENGTH Up to 13 m / 40 ft

DORSAL FIN Small, triangular and very far back on body

APPEARANCE Dark brown with irregular white patches on the belly; long rostrum (beak) and bulging forehead



Cuvier's beaked whale

Ziphius cavirostris

LENGTH Up to 7 m / 21 ft

DORSAL FIN More curved (falcate) like that of a dolphin

APPEARANCE Orange with small white patches and a stubby head



Stejneger's beaked whale

Mesoplodon stejnegeri

LENGTH Up to 5 m / 15 ft

DORSAL FIN Small, triangular and very far back on body

APPEARANCE Dark with light coloration around the lower jaw and eyes; prominent triangular tooth may be seen



Hubb's beaked whale

Mesoplodon carlhubbsi

LENGTH Up to 5 m / 15 ft

DORSAL FIN Small, triangular and very far back on body

APPEARANCE Dark with light coloration around both jaws and a white mound above the eye;

a triangular tooth may be seen





